

Inclusive Diffraction at HERA


Paul Newman, Birmingham University
representing the H1 and ZEUS Collaborations.



- Diffraction at HERA.
- The Diffractive Structure Function $F_2^{D(3)}$ at Low Q^2 .
- Rapidity Gaps Between High p_T Jets in Photoproduction.

Diffraction at HERA

At HERA, diffractive $\gamma^{(*)}p$ interactions can be studied ...




All five kinematic variables can be measured:

- $Q^2 \sim 0, |t| \sim 0$. \rightarrow similar to soft hadronic diffraction.
- Large Q^2 . \rightarrow pQCD at $\gamma^* IP$ vertex. γ^* 'probes' IP?
- Large $|t|$. \rightarrow IP itself calculable in pQCD?

... All regions interesting - transitions particularly revealing?

Diffraction of Virtual Photons, $\gamma^* p \rightarrow X p$




$$x_{\mathbb{P}} = \frac{q \cdot (p - p')}{q \cdot p} = x_{\mathbb{P}/p}$$

$$\beta = \frac{Q^2}{q \cdot (p - p')} = x_{q/\mathbb{P}}$$

$$(x = x_{\mathbb{P}} \beta)$$

Data presented as a Diffractive Structure Function ...

$$F_2^{D(3)}(\beta, Q^2, x_{\mathbb{P}})$$



New ZEUS BPC data ($0.2 < Q^2 < 0.7 \text{ GeV}^2$) start to fill transition to photoproduction.

Method 1: Decompose observed M_X distribution.

Method 2: Measure leading proton in LPS.


$x_{\mathbb{P}}$ Dependence of $F_2^{D(3)}$ at low Q^2

Diff. structure function $F_2^{D(3)}(\beta, Q^2, x_{\mathbb{P}})$ for $M_Y \lesssim 6$ GeV

By comparing M_X decomposition and LPS methods ...

P-dissociation contribution = 29 ± 15 (stat.) %

ZEUS 1996-97 PRELIMINARY (M_X method)



Fit to

$$F_2^{D(3)}(\beta, Q^2, x_{\mathbb{P}}) = A(\beta, Q^2) \cdot (1/x_{\mathbb{P}})^a$$

Consistent with
Regge factorisation
of $x_{\mathbb{P}}$ dependence

In Regge pole models, $a = \langle 2\alpha_{\mathbb{P}}(t) - 1 \rangle \dots$


$$\langle \alpha_{\mathbb{P}}(t) \rangle = 1.126 \pm 0.012 \text{ (stat.) } {}^{+0.027}_{-0.032} \text{ (syst.)}$$

Variation of Energy Dependence with Q^2

Expressed through Regge parameterisations ...

$$x_{\text{IP}} F_2^D \sim A(\beta, Q^2) x^{2-2\langle\alpha_{\text{IP}}(t)\rangle} \quad F_2 \sim B(Q^2) x^{1-\alpha_{\text{IP}}(0)}$$

Effective pomeron intercept $\alpha_{\text{IP}}(0)$, corrected for finite t ...





Also From LPS, $\alpha_{\text{IP}}(0) = 1.18 \pm 0.06$ (stat.) $^{+0.06}_{-0.09}$ (syst.)

At low Q^2 , diffractive and inclusive $\alpha_{\text{IP}}(0)$ compatible.



At higher Q^2 , diffractive and inclusive $\alpha_{\text{IP}}(0)$ incompatible.

(Energy dependences of diff & incl become more similar at high Q^2 .)

β / Q^2 dependence of $F_2^{D(3)}$




Rising scaling violations over large range of Q^2 up to large fractional momenta (β)



DGLAP analysis yields IP parton distributions dominated by "hard" gluons.

Q^2 Dependence of F_2^D

ZEUS 1995-1997 PRELIMINARY



\triangle M_X Prelim

\star LPS Prelim

\circ 94

— Sat'n ($q\bar{q} + q\bar{q}g$)

- - - Sat'n ($q\bar{q}$)

(94 and M_X data scaled to remove p-diss)

Clear transition at low Q^2 . Similar to that in total F_2 .

'Saturation' Model (Golec-Biernat & Wüsthoff)

Describes $Q^2 \rightarrow 0$ transition in F_2 and Q^2 dependence of diff, incl $\alpha_{\text{IP}}(0)$.

No new free parameters for diffraction!


Good agreement with high Q^2 diff data.

Diff'n of $\gamma^* \rightarrow q\bar{q}g$ dominates at low β .

Applicability of present $q\bar{q}g$ model questionable at low Q^2 .

Qualitative features of transition described.

ZEUS 1994




$x_{\text{IP}} = 0.0042$

β

Rapidity Gaps between Jets at $Q^2 = 0$

Parton level

Elastic parton-parton scattering
in Regge limit ($\hat{s} \gg \hat{t}$), yet
pQCD calculable (\hat{t} large)? ... BFKL?




Hadron level Classic experimental signature is rapidity gap
between high p_T jets. $|\hat{t}| \sim p_{t,jet}^2$

Complication: Remnant-remnant interactions produce
hadronic activity between jets?

New Measurement Method:

Require two central jets (inclusive k_t clustering algorithm):

$$E_t^1 > 6 \text{ GeV} \quad E_t^2 > 5 \text{ GeV} \quad \Delta\eta > 2.5$$



$$\mathbb{E}_t^{jets} = \sum E_t$$

for $\eta_f > \eta > \eta_b$.

outside two leading jets


Rapidity gap event if

$$\mathbb{E}_t^{jets} < E_t^{cut}$$

Vary E_t^{cut} to study effect
of spectator interactions.

Dependence of Gap Fraction on E_t^{cut}

f = Fraction of events with $\#_t^{jets} < E_t^{cut}$ ($\Delta\eta > 2.5$)



pQCD treatment of spectator interactions possible at large E_t^{cut} ?

Models with standard γp matrix elements and multiple interactions (HERWIG + JIMMY, PYTHIA) underestimate f

Models with high $|t|$ colour singlet exchange more successful

HERWIG + JIMMY + BFKL: LO BFKL calculation of $qq \rightarrow qq$ (q, g couplings. - $\alpha_s = 0.17$).

PYTHIA + ($\gamma \times 1200$): $qq \rightarrow qq$ through γ exchange.


(q coupling only - tuned to data).

Tuning of multiple interactions still required!

Dependence of Gap Fraction on $\Delta\eta$

Dependence on jet separation $\Delta\eta$ particularly sensitive to dynamics.

Measured for various E_t^{cut} , 1 GeV chosen here.



Clear signal above standard γp models, increases with $\Delta\eta$


Gap fraction at large $\Delta\eta$ significantly larger than Tevatron $p\bar{p}$.

Both models simulating colour singlet exchange describe data.

Dependence of Gap Fraction on x_p^{jets}

q, g composition of proton changes with x

x_p^{jets} dependence of f sensitive to q, g couplings of exchange.



Clear signal above standard γp models at all x_p^{jets} .

PYHTIA + γ differs in shape from HERWIG + BFKL.

Data favour some g coupling to exchange? - Improved statistics needed!

Summary

- New data in two previously unexplored regions ...
- F_2^D data for $0.2 < Q^2 < 0.7 \text{ GeV}^2$:
 - Effective \mathbb{P} intercept significantly larger than soft \mathbb{P} at high Q^2 .
 - Energy dependence of $\sigma^{\text{diff}} / \sigma^{\text{incl}}$ consistent with simple Regge prediction at low Q^2 .
 - Transition $Q^2 \rightarrow 0$ qualitatively similar to inclusive F_2
- Rapidity Gaps between Jets in Photoproduction:
 - Clear signal for colour singlet exchange at high $|t|$.
 - Good treatment of spectator interactions is crucial.
 - Sensitivity to q, g couplings of colour singlet exchange.